

**FLORIDA
MUSEUM**
OF NATURAL HISTORY®

BULLETIN

ISLAND LISTS OF WEST INDIAN AMPHIBIANS AND REPTILES

Robert Powell and Robert W. Henderson, Editors

Vol. 51, No. 2, pp. 85–166

2012

UNIVERSITY OF FLORIDA

GAINESVILLE

PUERTO RICO AND THE VIRGIN ISLANDS

Gregory C. Mayer¹

"The multitude of other islands in the group will well repay the visit of the zoologist, since the life of this section of the West Indies cannot be said to be known until all the cays and islets have been investigated" (Wetmore 1927).

Puerto Rico and the Virgin Islands comprise over 150 islands for which herpetological records exist. These islands are spread over five island banks: the Mona Bank, the Monito Bank, the Desecheo Bank, the Puerto Rico Bank (also known as Greater Puerto Rico: Schmidt 1928), and the St. Croix Bank (Fig. 8). The Puerto Rico Bank extends from the Mona Passage in the west to the Anegada Passage in the east. On the north, it is bordered by the Puerto Rican Trench, and to the south deep water separates it from St. Croix. The islands on this bank are Puerto Rico and adjacent islets, the Passage Islands (Vieques and Culebra and associated islets), the U.S. Virgin Islands (except St. Croix), and the British Virgin Islands. All of these islands were united into a single land mass during the low sea level stands associated with the Pleistocene glacial maxima, when sea level stood approximately 120 m below its present level (Heatwole & MacKenzie 1967, Fairbanks 1989). Because banks set the limits to recent over-land dispersal, they are key entities in herpetogeographic analyses (Barbour & Shreve 1935). The islands of the other banks are separated from the Puerto Rico Bank by water sufficiently deep to have precluded their being joined to Greater Puerto Rico by any glacial sea level drop. The Mona Bank, 60 km west of Puerto Rico, has a single island, Isla Mona. Two kilometers to the northwest, separated by water over 220 m deep, is the Monito Bank, with its single island, Isla Monito. The Desecheo Bank, 20 km west of Puerto Rico, also has but one island, Isla Desecheo. The fauna of these three islands is derived, in whole

or in part, from that of Puerto Rico, to which they belong politically. The St. Croix Bank, 40 km south of St. Thomas (on the Puerto Rico Bank), consists of St. Croix and four small islets. Its fauna shows considerable endemism, but also shares several forms with the Puerto Rico Bank.

About 100 species of reptiles and amphibians have been recorded from the area covered, of which more than 20 have been introduced. The introduced populations of *Cyclura pinguis* are deliberate translocations within the probable former range of this endangered species (Lazell 2005).

The following lists were compiled starting with the distributional checklist in Mayer (1989), to which I added the results of more recent fieldwork, data from the examination of museum specimens, and literature records. In addition to the general works cited in the introduction, the works of Stejneger (1904), Schmidt (1928), Grant (e.g., 1937), Philibosian and Yntema (1976, 1978), Rivero (1978, 1998), Heatwole et al. (1981), MacLean (1982), and Lazell (1983) are fundamental for Puerto Rican/Virgin Island herpetology. More recent works that have made useful compilations are Ovaska et al. (2000), Platenberg (2007), Perry and Gerber (2011), Platenberg and Boulon (2011), and Powell et al. (2011). Distributions within the large island of Puerto Rico are addressed in Rivero (1978, 1998), Schwartz and Henderson (1991), and Hedges (2012).

For areas and locations, the chief sources were McGuire (1925), Lazell (1983), the U.S. Geological Survey Geographic Names Information System (www.geonames.usgs.gov), the Montana State University Environmental Statistics Group (www.esg.montana.edu), and the National Geospatial-Intelligence Agency (<http://earth-info.nga.mil/gns/html/index.html>), supplemented by Google

¹Department of Biological Sciences, University of Wisconsin-Parkside, Kenosha, Wisconsin 53141 <mayerg@uwp.edu>




Figure 8. Map of greater Puerto Rico and adjacent island banks, including the Virgin Islands.

Earth® (www.earth.google.com) and Berkeley-Mapper (berkeleymapper.berkeley.edu). Coordinates for the BVI are less precise, as the NGA data are rounded to minutes. Many variant names exist for many of the islands, and in deciding which names to use, I have considered official names, navigational charts, local usage, and usage in herpetological and other scientific literature. I have not followed any one of these exclusively.

ACKNOWLEDGEMENTS

My primary debt is to James Lazell, who accompanied me through much of the field work, and provided support via The Conservation Agency. The Jarecki family, E.M. Azevedo, D. Ford, R.Z. Jenkins, W.P. and E. MacLean, N.R. Mayer, R. Thomas, and J. Walsh assisted in numerous ways, as did K. Tighe and J. Jacobs (National Museum of Natural History) and J. Rosado (Museum of Comparative Zoology) during the most recent museum work. James Lazell, Gad Perry, Kristiina Ovaska, and Renata J. Platenberg provided recent records.

MONA BANK

ISLA MONA (18.08°/-67.89°) (55.81 km²)
Eleutherodactylus monensis
Rhinella marina (I)

Chelonoidis monensis (F)

Ameiva alboguttata
Anolis monensis
Cyclura stejnegeri
Hemidactylus mabouia (I?)
Sphaerodactylus monensis
Spondylurus monae
Borikenophis variegatus
Epicrates monensis
Typhlops monensis

MONITO BANK

ISLA MONITO (18.16°/-67.95°) (0.16 km²)
Anolis monensis
Sphaerodactylus micropithecus
Spondylurus monitae

DESECHEO BANK

ISLA DESECHEO (18.39°/-67.48°) (1.46 km²)
Ameiva exsul
Anolis desechensis
Sphaerodactylus levinsi
Spondylurus nitidus
Borikenophis portoricensis ssp.

PUERTO RICO BANK

PUERTO RICO (18.25°/-66.50°) (8,674 km²)
Eleutherodactylus antillensis
Eleutherodactylus brittoni
Eleutherodactylus cochranae
Eleutherodactylus gryllus
Eleutherodactylus hedricki
Eleutherodactylus jasperi
Eleutherodactylus juanriveroi

Eleutherodactylus karlschmidti
Eleutherodactylus locustus
Eleutherodactylus portoricensis
Eleutherodactylus richmondi
Eleutherodactylus unicolor
Eleutherodactylus wightmanae
Hyla cinerea (I)
Leptodactylus albilabris
Leptodactylus fallax (W)
Lithobates grylio (I)
Lithobates catesbeianus (I)
Osteopilus septentrionalis (I)
Peltophryne lemur
Rhinella marina (I)
Scinax ruber (I)
Trachemys scripta elegans (I)
Trachemys stejnegeri stejnegeri
Ameiva exsul
Ameiva wetmorei
Amphisbaena bakeri
Amphisbaena caeca
Amphisbaena schmidti
Amphisbaena xera
Anolis cooki
Anolis cristatellus cristatellus
Anolis cristatellus cristatellus x A. c. wileyae
Anolis cuvieri
Anolis evermanni
Anolis gundlachi
Anolis krugi
Anolis occultus
Anolis poncensis
Anolis pulchellus
Anolis stratulus
Cyclura nubila (I)
Cyclura pinguis (F*)
Diploglossus pleii
Hemidactylus haitianus
Hemidactylus mabouia (I?)
Hemidactylus turcicus (W)
Iguana delicatissima (W)
Iguana iguana (I)
Leiocephalus etheridgei (F)
Leiocephalus partitus (F)
Phyllodactylus wirshingi
Scincella lateralis (I)
Sphaerodactylus gaigeae
Sphaerodactylus klauberi
Sphaerodactylus macrolepis ateles
Sphaerodactylus macrolepis grandisquamis
Sphaerodactylus macrolepis guarionex
Sphaerodactylus macrolepis mimetes
Sphaerodactylus macrolepis phoberus
Sphaerodactylus macrolepis spanius
Sphaerodactylus nicholsi
Sphaerodactylus roosevelti

Sphaerodactylus townsendi
Spondylurus nitidus
Boa constrictor ssp. (I)
Borikenophis portoricensis portoricensis
Borikenophis portoricensis prymnus
Epicrates inornatus
Epicrates granti
Eunectes notaeus (W)
Magliophis stahli
Magliophis exiguum subspadix
Python bivittatus (W)
Python regius (W)
Python reticulatus (W)
Python sebae (W)
Typhlops granti
Typhlops hypomethes
Typhlops platycephalus
Typhlops rostellatus
Caiman crocodilus (I)
CAYS OFF THE NORTHERN COAST OF PUERTO RICO
Isla de Cabras (= Leper Island) (18.47°/-66.14°)
Ameiva exsul
Anolis cristatellus cristatellus
Anolis pulchellus
Punta Salinas Island (18.47°/-66.18°)
Anolis cristatellus cristatellus
Anolis pulchellus
CAYS OFF THE SOUTHERN COAST OF PUERTO RICO
Caja de Muertos (17.89°/-66.52°)
Ameiva exsul
Ameiva wetmorei
Amphisbaena xera
Anolis cooki
Anolis cristatellus cristatellus
Anolis pulchellus
Phyllodactylus wirshingi
Sphaerodactylus roosevelti
Sphaerodactylus townsendi
Borikenophis portoricensis prymnus
Typhlops granti
Typhlops sp.
Cayos de Barca (= Cayos de la Barca) (17.92°/-66.24°)
Anolis cristatellus cristatellus
Cayo Cardona (17.96°/-66.63°)
Ameiva exsul
Anolis cristatellus cristatellus
Cayo Don Luis (17.94°/-66.97°)
Anolis cooki
Cayo Punta Aguila (17.95°/-67.21°)
Anolis cooki
Cayos de Ratones (near Jobos) (17.93°/-66.29°)
Anolis cristatellus cristatellus

| | |
|---|--|
| Cayo Turramote (17.94°/-67.02°) | Cayo Palominitos (18.34°/-65.57°) |
| <i>Sphaerodactylus nicholsi</i> (W) | <i>Eleutherodactylus cochranae</i> (W) |
| Isla Cueva (17.96°/-67.08°) | <i>Ameiva exsul</i> |
| <i>Hemidactylus mabouia</i> (I?) | <i>Anolis cristatellus wileyae</i> |
| Isla Magueyes (17.97°/-67.04°) | <i>Anolis pulchellus</i> (I, E) |
| <i>Rhinella marina</i> (I) | <i>Anolis stratulus</i> (W) |
| <i>Ameiva wetmorei</i> | <i>Typhlops</i> sp. |
| <i>Anolis cristatellus cristatellus</i> | |
| <i>Anolis poncensis</i> | |
| <i>Cyclura nubila</i> (I) | |
| <i>Iguana iguana</i> (I) | |
| <i>Sphaerodactylus nicholsi</i> | |
| Isla Mata Seca (17.96°/-67.01°) | Cayo Palominos (18.35°/-65.57°) |
| <i>Anolis cristatellus cristatellus</i> | <i>Ameiva exsul</i> |
| Isla Morrillito (= Platillo) (17.88°/-66.53°) | <i>Anolis cristatellus wileyae</i> |
| <i>Ameiva exsul</i> | |
| <i>Ameiva wetmorei</i> | |
| <i>Anolis cristatellus cristatellus</i> | |
| <i>Sphaerodactylus townsendi</i> | |
| <i>Borikenophis portoricensis prymnus</i> | |
| Isla Romero III (17.95°/-66.98°) | Cayo Piñeritos (18.24°/-65.59°) |
| <i>Anolis cristatellus cristatellus</i> | <i>Anolis cristatellus wileyae</i> |
| Long Island (= Long Key) (17.96°/-66.99°) | Cayo Santiago (18.16°/-65.73°) |
| <i>Anolis cristatellus cristatellus</i> | <i>Leptodactylus albilabris</i> |
| Spiny Butte (17.96°/-66.99°) | <i>Rhinella marina</i> (I) |
| <i>Ameiva exsul</i> | <i>Ameiva exsul</i> |
| <i>Anolis cristatellus cristatellus</i> | <i>Anolis cristatellus cristatellus</i> |
| <i>Sphaerodactylus nicholsi</i> | <i>Anolis pulchellus</i> |
| CAYS OFF THE EASTERN COAST OF PUERTO RICO | <i>Anolis stratulus</i> |
| Cabeza de Perro (= North Cabras Island) (18.25°/-65.58°) | <i>Sphaerodactylus gaigeae</i> |
| <i>Ameiva exsul</i> | <i>Sphaerodactylus macrolepis grandisquamis</i> |
| <i>Anolis cristatellus cristatellus</i> | <i>Borikenophis portoricensis portoricensis</i> |
| <i>Anolis pulchellus</i> | <i>Magliophis stahli</i> |
| <i>Sphaerodactylus macrolepis grandisquamis</i> | |
| <i>Borikenophis portoricensis</i> ssp. | |
| Cayo Ahogado (18.32°/-65.62°) | Isla Cabras (18.21°/-65.60°) |
| <i>Ameiva exsul</i> (W) | <i>Ameiva exsul</i> |
| <i>Anolis cristatellus cristatellus</i> (W) | <i>Anolis cristatellus</i> ssp. |
| <i>Anolis pulchellus</i> (W) | <i>Sphaerodactylus townsendi</i> |
| Cayo Algodones (18.19°/-65.68°) | Isla Piñeros (18.25°/-65.59°) |
| <i>Leptodactylus albilabris</i> | <i>Eleutherodactylus antillensis</i> |
| <i>Rhinella marina</i> (I) | <i>Leptodactylus albilabris</i> |
| <i>Ameiva exsul</i> | <i>Ameiva exsul</i> |
| <i>Anolis cristatellus cristatellus</i> x <i>A. c. wileyae</i> | <i>Amphisbaena caeca</i> |
| <i>Anolis pulchellus</i> | <i>Anolis cristatellus wileyae</i> |
| <i>Sphaerodactylus macrolepis grandisquamis</i> | <i>Anolis pulchellus</i> |
| Cayo Batata (18.12°/-65.77°) | <i>Anolis stratulus</i> |
| <i>Ameiva exsul</i> | <i>Sphaerodactylus gaigeae</i> |
| <i>Anolis cristatellus cristatellus</i> | <i>Sphaerodactylus macrolepis stibarus</i> |
| <i>Anolis pulchellus</i> | <i>Sphaerodactylus townsendi</i> |
| <i>Sphaerodactylus macrolepis grandisquamis</i> | <i>Borikenophis portoricensis</i> ssp. |
| | <i>Typhlops hypomethes</i> |
| | |
| | Isla de Ramos (18.31°/-65.61°) |
| | <i>Leptodactylus albilabris</i> |
| | <i>Ameiva exsul</i> |
| | <i>Anolis cristatellus cristatellus</i> x <i>A. c. wileyae</i> |
| | <i>Anolis pulchellus</i> |
| | <i>Sphaerodactylus townsendi</i> |
| | |
| | Isleta Marina (= Cayo Zancudo) (18.34°/-65.62°) |
| | <i>Ameiva exsul</i> |

Anolis cristatellus cristatellus x *A. c. wileyae*
Anolis pulchellus
Levins Rock (18.18°/-65.71°)
Ameiva exsul (W)
CAYOS DE LA CORDILLERA
La Blanquilla (18.37°/-65.55°)
Ameiva exsul
Anolis cristatellus wileyae
Sphaerodactylus townsendi
Booby Hatch Cay (= Booby Rock) (18.37°/-65.55°)
Anolis cristatellus wileyae
Sphaerodactylus townsendi
Cayo Diablo (= Cayo La Llave) (18.36°/-65.53°)
Ameiva exsul
Anolis cristatellus wileyae
Epicrates granti
Typhlops richardii
Cayo Icacos (= Hicacos) (18.39°/-65.59°)
Leptodactylus albilabris
Ameiva exsul
Anolis cristatellus wileyae
Anolis pulchellus
Iguana iguana (I)
Sphaerodactylus townsendi
Spondylurus nitidus
Cayo Lobos (18.38°/-65.57°)
Ameiva exsul
Anolis cristatellus wileyae
Anolis pulchellus
Sphaerodactylus townsendi
Cayo Ratones (off Fajardo) (18.38°/-65.58°)
Ameiva exsul
Anolis cristatellus wileyae
Anolis pulchellus
Sphaerodactylus townsendi
Epicrates granti (I)
Grant Rock (18.37°/-65.55°)
Anolis cristatellus wileyae
Sphaerodactylus townsendi
Konyoki (18.38°/-65.58°)
Ameiva exsul
Anolis cristatellus wileyae
Sphaerodactylus townsendi
MacKenzie Key (18.37°/-65.55°)
Anolis cristatellus wileyae
Sphaerodactylus townsendi
Pressick Rock (18.38°/-65.57°)
Sphaerodactylus townsendi
Surprise Key (18.37°/-65.55°)
Anolis cristatellus wileyae

Sphaerodactylus townsendi
Typhlops richardii
VIEQUES (= Bieque = Crab Island) (18.12°/-65.42°) (137.66 km²)
Eleutherodactylus antillensis
Eleutherodactylus cochranae
Eleutherodactylus coqui (?)
Leptodactylus albilabris
Lithobates catesbeianus (I)
Rhinella marina (I)
Trachemys stejnegeri stejnegeri (?)
Ameiva exsul
Anolis cristatellus wileyae
Anolis pulchellus
Anolis roosevelti (PE)
Anolis stratulus
Hemidactylus mabouia (?)
Sphaerodactylus gaigeae
Sphaerodactylus macrolepis inigoi
Sphaerodactylus roosevelti
Sphaerodactylus townsendi
Spondylurus nitidus (PX)
Borikenophis portoricensis aphantus
Typhlops hypomethes
Typhlops platycephalus
Caiman crocodilus (W)
Cayo de Afuera (= Cayo Real) (18.09°/-65.47°)
Ameiva exsul
Amphisbaena caeca
Sphaerodactylus macrolepis inigoi
Sphaerodactylus townsendi
Cayo de Tierra (18.09°/-65.47°)
Ameiva exsul
Hemidactylus mabouia (?)
Sphaerodactylus macrolepis inigoi
Sphaerodactylus townsendi
Typhlops sp.
Isla Chiva (18.11°/-65.38°)
Anolis cristatellus wileyae
Sphaerodactylus townsendi
CULEBRA (18.31°/-65.28°) (27.28 km²)
Eleutherodactylus antillensis
Eleutherodactylus cochranae
Eleutherodactylus coqui (I)
Leptodactylus albilabris
Rhinella marina (I)
Trachemys stejnegeri stejnegeri (?)
Ameiva exsul
Anolis cristatellus wileyae
Anolis pulchellus
Anolis roosevelti (PE)
Anolis stratulus
Hemidactylus mabouia (?)
Iguana iguana (I)

| | |
|---|--|
| <i>Sphaerodactylus macrolepis macrolepis</i> | <i>Chelonoidis carbonaria</i> (I?) |
| <i>Sphaerodactylus townsendi</i> | <i>Trachemys scripta elegans</i> (I) |
| <i>Spondylurus culebrae</i> | <i>Ameiva exsul</i> |
| <i>Spondylurus nitidus</i> | <i>Amphisbaena fenestrata</i> |
| <i>Borikenophis portoricensis richardi</i> | <i>Anolis leachii</i> (I, E?) |
| <i>Epicrates granti</i> | <i>Anolis cristatellus wileyae</i> |
| <i>Magliophis exiguis exiguis</i> | <i>Anolis pulchellus</i> |
| <i>Typhlops richardii</i> | <i>Anolis stratulus</i> |
| Cayo Botella (= Heatwole Island) (18.32°/-65.24°) | <i>Cyclura pinguis</i> (F*) |
| <i>Anolis cristatellus wileyae</i> | <i>Gymnophthalmus underwoodi</i> (I?) |
| <i>Sphaerodactylus macrolepis macrolepis</i> | <i>Hemidactylus mabouia</i> (I?) |
| Cayo Geniqui East (= Palada Cays) (18.34°/-65.23°) | <i>Iguana iguana</i> (I) |
| <i>Sphaerodactylus macrolepis macrolepis</i> | <i>Sphaerodactylus macrolepis macrolepis</i> |
| Cayo Luis Peña (= Southwest Key) (18.31°/-65.33°) | <i>Spondylurus semitaeniatus</i> (PX) |
| <i>Anolis cristatellus wileyae</i> | <i>Spondylurus sloanii</i> (PX) |
| <i>Anolis pulchellus</i> | <i>Spondylurus spilonotus</i> (PE) |
| <i>Sphaerodactylus macrolepis macrolepis</i> | <i>Thecadactylus rapicauda</i> (PX) |
| <i>Spondylurus nitidus</i> | <i>Boa constrictor</i> ssp. (W) |
| <i>Borikenophis portoricensis richardi</i> | <i>Borikenophis portoricensis richardi</i> (PX, I) |
| <i>Typhlops richardii</i> | <i>Epicrates granti</i> |
| Cayo Norte (18.34°/-65.26°) | <i>Magliophis exiguis exiguis</i> |
| <i>Ameiva exsul</i> | <i>Pantherophis guttatus</i> (I) |
| <i>Anolis cristatellus wileyae</i> | <i>Python regius</i> (W) |
| <i>Anolis pulchellus</i> | <i>Typhlops richardii</i> |
| <i>Sphaerodactylus macrolepis macrolepis</i> | Big Flat Cay (18.32°/-64.99°) (0.012 km ²) |
| <i>Spondylurus nitidus</i> | <i>Ameiva exsul</i> |
| <i>Typhlops richardii</i> | <i>Anolis cristatellus wileyae</i> |
| Cayo Pirata (= Chicken Island) (18.31°/-65.30°) | <i>Sphaerodactylus macrolepis macrolepis</i> |
| <i>Anolis cristatellus wileyae</i> | Bovoni Cay (18.31°/-64.87°) (0.20 km ²) |
| <i>Sphaerodactylus macrolepis macrolepis</i> | <i>Eleutherodactylus cochranae</i> |
| Culebrita (18.31°/-65.23°) | <i>Ameiva exsul</i> |
| <i>Ameiva exsul</i> | <i>Anolis cristatellus wileyae</i> |
| <i>Anolis cristatellus wileyae</i> | <i>Anolis stratulus</i> |
| <i>Sphaerodactylus macrolepis macrolepis</i> | <i>Hemidactylus mabouia</i> (I?) |
| <i>Spondylurus culebrae</i> | <i>Sphaerodactylus macrolepis macrolepis</i> |
| <i>Typhlops richardii</i> | Buck Island (western and larger of the Capella Islands) (18.28°/-64.89°) (0.17 km ²) |
| Isla Pela (= Bare Cactus Island) (18.30°/-65.25°) | <i>Anolis cristatellus wileyae</i> |
| <i>Anolis cristatellus wileyae</i> | <i>Sphaerodactylus macrolepis macrolepis</i> |
| <i>Sphaerodactylus macrolepis macrolepis</i> | <i>Spondylurus semitaeniatus</i> |
| Pelaita (= Villa del Mar) (18.30°/-65.25°) | <i>Spondylurus sloanii</i> |
| <i>Anolis cristatellus wileyae</i> | <i>Borikenophis portoricensis nicholsi</i> |
| <i>Sphaerodactylus macrolepis macrolepis</i> | Capella Island (= Eastern Island = unnamed island, eastern and smaller of the Capella Islands) (18.28°/-64.89°) (0.09 km ²) |
| ST. THOMAS (18.35°/-64.94°) (70.23 km ²) | <i>Anolis cristatellus wileyae</i> |
| <i>Eleutherodactylus antillensis</i> | <i>Spondylurus semitaeniatus</i> |
| <i>Eleutherodactylus cochranae</i> | <i>Spondylurus sloanii</i> |
| <i>Eleutherodactylus coqui</i> (I) | Cas Cay (18.31°/-64.87°) (0.06 km ²) |
| <i>Eleutherodactylus latus</i> | <i>Ameiva exsul</i> |
| <i>Eleutherodactylus schwartzi</i> (?) | <i>Anolis cristatellus wileyae</i> |
| <i>Leptodactylus albilibris</i> | <i>Anolis stratulus</i> |
| <i>Osteopilus septentrionalis</i> (I) | <i>Iguana iguana</i> (I) |
| <i>Rhinella marina</i> (I) | <i>Sphaerodactylus macrolepis macrolepis</i> |

Cockroach Island (18.40°/-65.06°) (0.08 km²)

Anolis cristatellus wileyae
Sphaerodactylus macrolepis macrolepis
Borikenophis portoricensis richardi

Congo Cay (18.37°/-64.80°) (0.10 km²)

Anolis cristatellus wileyae
Anolis stratulus
Sphaerodactylus macrolepis macrolepis
Borikenophis portoricensis richardi
Typhlops richardii

Current Rock (18.31°/-64.83°) (0.002 km²)

Anolis cristatellus wileyae

Dog Island (18.30°/-64.82°) (0.05 km²)

Ameiva exsul

Anolis cristatellus wileyae
Sphaerodactylus macrolepis macrolepis
Borikenophis portoricensis richardi

Dutcheap Cay (= Dutchman's Cap) (18.38°/-65.06°) (0.13 km²)

Ameiva exsul

Anolis cristatellus wileyae
Sphaerodactylus macrolepis macrolepis
Borikenophis portoricensis richardi

Fish Cay (18.31°/-64.83°) (0.001 km²)

Anolis cristatellus wileyae

Hemidactylus mabouia (I?)

French Cap Cay (= Frenchman's Cap) (18.23°/-64.85°) (0.04 km²)

Sphaerodactylus townsendi

Grass Cay (18.36°/-64.83°) (0.20 km²)

Anolis cristatellus wileyae

Sphaerodactylus macrolepis macrolepis
Borikenophis portoricensis richardi

Great St. James Island (18.31°/-64.83°) (0.63 km²)

Ameiva exsul

Amphisbaena fenestrata

Anolis cristatellus wileyae

Anolis pulchellus

Anolis stratulus

Sphaerodactylus macrolepis macrolepis

Spondylurus sp.

Borikenophis portoricensis richardi

Hans Lollik Island (18.40°/-64.91°) (1.98 km²)

Ameiva exsul

Anolis cristatellus wileyae

Anolis pulchellus

Anolis stratulus

Sphaerodactylus macrolepis macrolepis

Borikenophis portoricensis richardi

Hassel Island (18.33°/-64.94°) (0.56 km²)

Eleutherodactylus cochranae

Eleutherodactylus latus

Osteopilus septentrionalis (I)

Chelonoidis carbonaria (I?)

Ameiva exsul

Anolis cristatellus wileyae

Anolis pulchellus

Anolis stratulus

Hemidactylus mabouia (I?)

Iguana iguana (I)

Sphaerodactylus macrolepis macrolepis

Magliophis exiguum *exiguus*

Inner Brass Island (18.38°/-64.97°) (0.52 km²)

Osteopilus septentrionalis (I)

Ameiva exsul

Anolis cristatellus wileyae

Anolis stratulus

Sphaerodactylus macrolepis macrolepis (?)

Borikenophis portoricensis ssp. (?)

Little Hans Lollik Island (18.41°/-64.91°) (0.41 km²)

Ameiva exsul

Anolis cristatellus wileyae

Sphaerodactylus macrolepis macrolepis

Typhlops richardii

Little St. James Island (18.30°/-64.83°) (0.28 km²)

Eleutherodactylus sp. (W)

Osteopilus septentrionalis (I)

Ameiva exsul

Anolis cristatellus wileyae

Anolis pulchellus

Anolis stratulus

Iguana iguana (I)

Sphaerodactylus macrolepis macrolepis

Borikenophis portoricensis richardi

Pantherophis guttatus (W)

Lovango Cay (18.36°/-64.81°) (0.48 km²)

Chelonoidis carbonaria (I?)

Ameiva exsul

Anolis cristatellus wileyae

Anolis pulchellus

Anolis stratulus

Hemidactylus mabouia (I?)

Borikenophis portoricensis richardi

Mingo Cay (18.36°/-64.82°) (0.20 km²)

Ameiva exsul

Anolis cristatellus wileyae

Anolis pulchellus

Anolis stratulus

Sphaerodactylus macrolepis macrolepis

Borikenophis portoricensis richardi

Outer Brass Island (18.39°/-64.97°) (0.44 km²)

Ameiva exsul

Anolis cristatellus wileyae

Anolis pulchellus

Anolis stratulus

Sphaerodactylus macrolepis macrolepis

Borikenophis portoricensis richardi

| | |
|--|--|
| Patricia Cay (18.31°/-64.87°) (0.14 km ²) | West Cay (18.36°/-65.05°) (0.16 km ²) |
| <i>Ameiva exsul</i> | <i>Ameiva exsul</i> |
| <i>Anolis cristatellus wileyae</i> | <i>Anolis cristatellus wileyae</i> |
| <i>Anolis stratulus</i> | <i>Anolis stratulus</i> |
| <i>Hemidactylus mabouia</i> (?) | <i>Sphaerodactylus macrolepis macrolepis</i> |
| <i>Iguana iguana</i> (I) | ST. JOHN (18.34°/-64.75°) (49.67 km ²) |
| <i>Sphaerodactylus macrolepis macrolepis</i> | <i>Eleutherodactylus antillensis</i> |
| Rotto Cay (18.31°/-64.86°) (0.008 km ²) | <i>Eleutherodactylus cochranae</i> |
| <i>Ameiva exsul</i> | <i>Eleutherodactylus coqui</i> (I) |
| <i>Anolis cristatellus wileyae</i> | <i>Eleutherodactylus latus</i> |
| <i>Sphaerodactylus macrolepis macrolepis</i> | <i>Eleutherodactylus schwartzi</i> (PX) |
| Saba Island (= Little Saba Island) (18.31°/-65.00°) (0.12 km ²) | <i>Leptodactylus albilabris</i> |
| <i>Ameiva exsul</i> | <i>Osteopilus septentrionalis</i> (I) |
| <i>Anolis cristatellus wileyae</i> | <i>Chelonoidis carbonaria</i> (?) |
| <i>Anolis stratulus</i> | <i>Ameiva exsul</i> |
| <i>Sphaerodactylus macrolepis macrolepis</i> | <i>Amphisbaena fenestrata</i> |
| <i>Spondylurus sloanii</i> | <i>Anolis cristatellus wileyae</i> |
| <i>Borikenophis portoricensis richardi</i> | <i>Anolis pulchellus</i> |
| Sail Rock (18.29°/-65.10°) (0.007 km ²) | <i>Anolis roosevelti</i> (PE) |
| <i>Sphaerodactylus macrolepis macrolepis</i> | <i>Anolis stratulus</i> |
| Salt Cay (18.36°/-65.05°) (0.23 km ²) | <i>Hemidactylus mabouia</i> (?) |
| <i>Ameiva exsul</i> | <i>Iguana iguana</i> (I) |
| <i>Anolis cristatellus wileyae</i> | <i>Sphaerodactylus macrolepis macrolepis</i> |
| <i>Sphaerodactylus macrolepis macrolepis</i> | <i>Spondylurus spilonotus</i> (PE) |
| <i>Spondylurus</i> sp. | <i>Borikenophis portoricensis richardi</i> (PX) |
| Savana Island (18.34°/-65.08°) (0.70 km ²) | <i>Magliophis exiguis</i> <i>exiguus</i> |
| <i>Ameiva exsul</i> | <i>Typhlops richardii</i> |
| <i>Anolis cristatellus wileyae</i> | Cinnamon Cay (18.36°/-64.76°) (0.004 km ²) |
| <i>Anolis stratulus</i> | <i>Anolis cristatellus wileyae</i> |
| <i>Sphaerodactylus macrolepis macrolepis</i> | <i>Sphaerodactylus macrolepis macrolepis</i> |
| <i>Borikenophis portoricensis richardi</i> | Cocoloba Cay (18.32°/-64.76°) (0.004 km ²) |
| Shark Island (18.34°/-64.84°) (0.005 km ²) | <i>Anolis cristatellus wileyae</i> |
| <i>Anolis cristatellus wileyae</i> | <i>Sphaerodactylus macrolepis macrolepis</i> |
| Thatch Cay (18.36°/-64.86°) (0.96 km ²) | Flanagan Island (18.33°/-64.65°) (0.09 km ²) |
| <i>Eleutherodactylus cochranae</i> | <i>Ameiva exsul</i> |
| <i>Ameiva exsul</i> | <i>Anolis cristatellus wileyae</i> |
| <i>Anolis cristatellus wileyae</i> | <i>Anolis stratulus</i> |
| <i>Anolis stratulus</i> | <i>Sphaerodactylus macrolepis macrolepis</i> |
| <i>Sphaerodactylus macrolepis macrolepis</i> | <i>Spondylurus</i> sp. |
| <i>Borikenophis portoricensis richardi</i> | <i>Typhlops richardii</i> |
| Turtledove Cay (18.31°/-65.00) (.02 km ²) | Henley Cay (one of the Durloe Cays) (18.35°/-64.79°) (0.05 km ²) |
| <i>Anolis cristatellus wileyae</i> | <i>Anolis cristatellus wileyae</i> |
| <i>Sphaerodactylus macrolepis macrolepis</i> | <i>Hemidactylus mabouia</i> (?) |
| Water Island (18.32°/-64.95°) (1.99 km ²) | <i>Sphaerodactylus macrolepis macrolepis</i> |
| <i>Chelonoidis carbonaria</i> (?) | Leduck Island (18.32°/-64.69°) (0.05 km ²) |
| <i>Ameiva exsul</i> | <i>Ameiva exsul</i> |
| <i>Anolis cristatellus wileyae</i> | <i>Anolis cristatellus wileyae</i> |
| <i>Anolis stratulus</i> | <i>Anolis stratulus</i> |
| <i>Hemidactylus mabouia</i> (?) | Ramgoat Cay (18.36°/-64.79°) (0.011 km ²) |
| <i>Iguana iguana</i> (I) | <i>Anolis cristatellus wileyae</i> |
| <i>Sphaerodactylus macrolepis macrolepis</i> | Rata Cay (18.36°/-64.80°) (0.002 km ²) |
| <i>Spondylurus sloanii</i> (PX) | <i>Anolis cristatellus wileyae</i> |
| <i>Borikenophis portoricensis richardi</i> | |

Steven Cay (18.33°/-64.81°) (0.008 km²)

Anolis cristatellus wileyae
Anolis stratulus
Iguana iguana (I)
Sphaerodactylus macrolepis macrolepis
Epicrates granti (I)

Trunk Cay (18.36°/-64.77°) (0.009 km²)

Anolis cristatellus wileyae
Anolis stratulus

Waterlemon Cay (18.37°/-64.72°) (0.003 km²)

Anolis cristatellus wileyae
Sphaerodactylus macrolepis macrolepis

Whistling Cay (18.37°/-64.75°) (0.08 km²)

Anolis cristatellus wileyae
Anolis stratulus
Iguana iguana (I)
Sphaerodactylus macrolepis macrolepis

JOST VAN DYKE (18.47°/-64.75°) (8.40 km²)

Eleutherodactylus antillensis
Eleutherodactylus latus (I?)
Eleutherodactylus schwartzi
Leptodactylus albilabris
Amphisbaena fenestrata
Anolis cristatellus wileyae
Anolis pulchellus
Anolis stratulus
Hemidactylus mabouia (I?)
Iguana iguana (W?)
Sphaerodactylus macrolepis macrolepis
Borikenophis portoricensis ssp. (PX)
Epicrates granti
Magliophis exiguus exiguus
Typhlops richardii

Great Tobago (18.45°/-64.80°) (0.87 km²)

Ameiva exsul
Anolis cristatellus wileyae
Hemidactylus mabouia (I?)
Sphaerodactylus macrolepis macrolepis

Green Cay (18.45°/-64.70°) (0.07 km²)

Anolis cristatellus wileyae
Sphaerodactylus macrolepis macrolepis
Borikenophis portoricensis ssp.

Little Jost van Dyke (18.45°/-64.72°) (0.63 km²)

Ameiva exsul
Anolis cristatellus wileyae
Anolis pulchellus
Sphaerodactylus macrolepis macrolepis
Borikenophis portoricensis ssp.
Typhlops richardii

Little Tobago (18.43°/-64.85°) (0.22 km²)

Anolis cristatellus wileyae
Sphaerodactylus macrolepis macrolepis

Spondylurus sloanii

Borikenophis portoricensis ssp.

Sandy Cay (18.43°/-64.72°) (0.06 km²)

Ameiva exsul
Anolis cristatellus wileyae
Hemidactylus mabouia (I?)
Sphaerodactylus macrolepis macrolepis

Sandy Spit (18.43°/-64.70°) (0.001 km²)

Anolis cristatellus wileyae
Sphaerodactylus macrolepis macrolepis

Watson Rock (= Cable Rock = Shark Rock) (18.44°/-64.83°)

Sphaerodactylus macrolepis macrolepis

TORTOLA (18.45°/-64.60°) (54.44 km²)

Eleutherodactylus antillensis
Eleutherodactylus cochranae
Eleutherodactylus johnstonei (I)
Eleutherodactylus schwartzzi
Leptodactylus albilabris
Osteopilus septentrionalis (I)
Chelonoidis carbonaria (I?)
Pseudemys nelsoni (I)

Trachemys scripta elegans (I)

Ameiva exsul
Ameiva sp. (I?)
Amphisbaena fenestrata
Anolis cristatellus wileyae
Anolis pulchellus

Anolis roosevelti (PE)
Anolis stratulus
Hemidactylus mabouia (I?)
Iguana iguana (I)

Sphaerodactylus macrolepis macrolepis
Sphaerodactylus parthenopion

Spondylurus semitaeniatus
Borikenophis portoricensis richardi
Epicrates granti
Magliophis exiguus exiguus
Pantherophis guttatus (W?)
Typhlops richardii

Beef Island (18.45°/-64.52°) (3.72 km²)

Eleutherodactylus antillensis
Eleutherodactylus schwartzzi
Leptodactylus albilabris
Osteopilus septentrionalis (I)

Ameiva exsul
Anolis cristatellus wileyae
Anolis pulchellus
Anolis stratulus

Hemidactylus mabouia (I?)
Sphaerodactylus macrolepis macrolepis
Borikenophis portoricensis anagadae
Magliophis exiguus exiguus
Typhlops richardii

| | |
|---|--|
| Bellamy Cay (18.43°/-64.53°) (0.007 km ²) | <i>Borikenophis portoricensis anegadae</i> |
| <i>Anolis cristatellus wileyae</i> | <i>Epicrates granti</i> (?) |
| <i>Hemidactylus mabouia</i> (I?) | <i>Magliophis exiguum exiguum</i> |
| <i>Sphaerodactylus macrolepis macrolepis</i> | <i>Typhlops richardii</i> |
| Buck Island (18.42°/-64.55°) (0.17 km ²) | Little Camanoe (18.45°/-64.53°) (0.16 km ²) |
| <i>Ameiva exsul</i> | <i>Ameiva exsul</i> |
| <i>Anolis cristatellus wileyae</i> | <i>Anolis cristatellus wileyae</i> |
| <i>Sphaerodactylus macrolepis macrolepis</i> | <i>Sphaerodactylus macrolepis macrolepis</i> |
| Frenchman's Cay (18.38°/-64.70°) (0.24 km ²) | Little Thatch Island (18.38°/-64.72°) (0.24 km ²) |
| <i>Eleutherodactylus antillensis</i> | <i>Eleutherodactylus schwartzi</i> (I) |
| <i>Eleutherodactylus schwartzii</i> | <i>Anolis cristatellus wileyae</i> |
| <i>Ameiva exsul</i> | <i>Anolis stratulus</i> |
| <i>Amphisbaena fenestrata</i> | <i>Cyclura pinguis</i> (I) |
| <i>Anolis cristatellus wileyae</i> | <i>Sphaerodactylus macrolepis macrolepis</i> |
| <i>Anolis pulchellus</i> | <i>Spondylurus semitaeniatus</i> |
| <i>Anolis stratulus</i> | Marina Cay (18.45°/-64.52°) (0.010 km ²) |
| <i>Hemidactylus mabouia</i> (I?) | <i>Ameiva exsul</i> |
| <i>Sphaerodactylus macrolepis macrolepis</i> | <i>Anolis cristatellus wileyae</i> |
| <i>Magliophis exiguum exiguum</i> | <i>Anolis stratulus</i> |
| Great Camanoe (18.48°/-64.53°) (3.37 km ²) | <i>Hemidactylus mabouia</i> (I?) |
| <i>Eleutherodactylus antillensis</i> | <i>Sphaerodactylus macrolepis macrolepis</i> |
| <i>Ameiva exsul</i> | <i>Typhlops richardii</i> |
| <i>Amphisbaena fenestrata</i> | Scrub Island (18.47°/-64.52°) (0.97 km ²) |
| <i>Anolis cristatellus wileyae</i> | <i>Ameiva exsul</i> |
| <i>Anolis pulchellus</i> | <i>Anolis cristatellus wileyae</i> |
| <i>Anolis stratulus</i> | <i>Anolis stratulus</i> |
| <i>Hemidactylus mabouia</i> (I?) | <i>Sphaerodactylus macrolepis macrolepis</i> |
| <i>Sphaerodactylus macrolepis macrolepis</i> | <i>Spondylurus sp.</i> |
| <i>Spondylurus semitaeniatus</i> | PETER ISLAND (18.37°/-64.58°) (4.29 km ²) |
| <i>Borikenophis portoricensis anegadae</i> | <i>Eleutherodactylus</i> sp. |
| <i>Epicrates granti</i> | <i>Osteopilus septentrionalis</i> (I) |
| <i>Magliophis exiguum exiguum</i> | <i>Chelonoidis carbonaria</i> (I?) |
| Great Thatch Island (18.38°/-64.72°) (1.23 km ²) | <i>Ameiva exsul</i> |
| <i>Eleutherodactylus antillensis</i> | <i>Anolis cristatellus wileyae</i> |
| <i>Eleutherodactylus cochranae</i> | <i>Anolis pulchellus</i> |
| <i>Amphisbaena fenestrata</i> | <i>Anolis stratulus</i> |
| <i>Anolis cristatellus wileyae</i> | <i>Hemidactylus mabouia</i> (I?) |
| <i>Anolis pulchellus</i> | <i>Iguana iguana</i> (I) |
| <i>Anolis stratulus</i> | <i>Sphaerodactylus macrolepis macrolepis</i> |
| <i>Sphaerodactylus macrolepis macrolepis</i> | <i>Spondylurus sloanii</i> |
| Guana Island (18.48°/-64.57°) (2.97 km ²) | <i>Borikenophis portoricensis richardi</i> |
| <i>Eleutherodactylus antillensis</i> | <i>Magliophis exiguum exiguum</i> |
| <i>Osteopilus septentrionalis</i> (W) | <i>Pantherophis guttatus</i> (W) |
| <i>Centrochelys sulcata</i> (I) | Carrot Rock (18.32°/-64.57°) (0.013 km ²) |
| <i>Chelonoidis carbonaria</i> (I) | <i>Anolis ernestwilliamsi</i> |
| <i>Ameiva exsul</i> | <i>Sphaerodactylus macrolepis macrolepis</i> |
| <i>Amphisbaena fenestrata</i> | <i>Spondylurus macleani</i> |
| <i>Anolis cristatellus wileyae</i> | Deadman's Chest (= Dead Chest) (18.37°/-64.57°) (0.14 km ²) |
| <i>Anolis pulchellus</i> | <i>Ameiva exsul</i> |
| <i>Anolis stratulus</i> | <i>Anolis cristatellus wileyae</i> |
| <i>Cyclura pinguis</i> (I) | <i>Sphaerodactylus macrolepis macrolepis</i> |
| <i>Hemidactylus mabouia</i> (I?) | <i>Spondylurus sp.</i> |
| <i>Sphaerodactylus macrolepis macrolepis</i> | |
| <i>Spondylurus semitaeniatus</i> | |

Norman Island (18.33° /- 64.62°) (2.57 km^2)
Ameiva exsul
Anolis cristatellus wileyae
Anolis pulchellus
Anolis stratulus
Cyclura pinguis (I)
Sphaerodactylus macrolepis macrolepis
Spondylurus sloanii
Borikenophis portoricensis ssp.
Typhlops richardii

Pelican Island (18.33° /- 64.62°) (0.03 km^2)
Anolis cristatellus wileyae
Sphaerodactylus macrolepis macrolepis

Skipper Cay (= Key Cay) (18.34° /- 64.60°) (0.011 km^2)
Anolis cristatellus wileyae

VIRGIN GORDA (18.50° /- 64.40°) (21.30 km^2)
Eleutherodactylus antillensis
Eleutherodactylus cochranae
Eleutherodactylus schwartzii
Osteopilus septentrionalis (I)
Peltophryne lemur (PX)
Chelonoidis carbonaria (I?)
Ameiva exsul
Amphisbaena fenestrata
Anolis cristatellus wileyae
Anolis pulchellus
Anolis stratulus
Hemidactylus mabouia (I?)
Iguana iguana (I)
Sphaerodactylus macrolepis macrolepis
Sphaerodactylus parthenopion
Spondylurus semitaeniatus
Borikenophis portoricensis anegadae
Magliophis exiguum
Typhlops naugus

Broken Jerusalem (18.40° /- 64.45°) (0.009 km^2)
Anolis cristatellus wileyae

Cistern Rock (18.38° /- 64.50°) (0.001 km^2)
Anolis cristatellus wileyae

Cooper Island (18.37° /- 64.50°) (1.38 km^2)
Ameiva exsul
Anolis cristatellus wileyae
Anolis stratulus
Hemidactylus mabouia (I?)
Sphaerodactylus macrolepis macrolepis
Borikenophis portoricensis ssp.

East Seal Dog (18.50° /- 64.43°) (0.008 km^2)
Anolis cristatellus wileyae
Sphaerodactylus macrolepis macrolepis

Eustatia Island (18.52° /- 64.37°) (0.11 km^2)
Ameiva exsul
Anolis cristatellus wileyae
Sphaerodactylus macrolepis macrolepis

Fallen Jerusalem (18.42° /- 64.45°) (0.12 km^2)
Anolis cristatellus wileyae
Anolis pulchellus
Anolis stratulus
Sphaerodactylus macrolepis macrolepis
Spondylurus semitaeniatus

George Dog (18.5° /- 64.45°) (0.15 km^2)
Ameiva exsul
Anolis cristatellus wileyae
Sphaerodactylus macrolepis macrolepis

Ginger Island (18.40° /- 64.47°) (1.05 km^2)
Ameiva exsul
Anolis cristatellus wileyae
Anolis stratulus
Sphaerodactylus macrolepis macrolepis
Spondylurus semitaeniatus
Borikenophis portoricensis ssp.

Great Dog (18.48° /- 64.45°) (0.33 km^2)
Eleutherodactylus schwartzii
Ameiva exsul
Anolis cristatellus wileyae
Sphaerodactylus macrolepis macrolepis
Typhlops sp.

Mosquito Island (= Mosquito Island) (18.50° /- 64.38°) (0.50 km^2)
Osteopilus septentrionalis (I)
Ameiva exsul
Anolis cristatellus wileyae
Anolis stratulus
Cyclura pinguis (I)
Hemidactylus mabouia (I?)
Sphaerodactylus macrolepis macrolepis
Sphaerodactylus parthenopion
Spondylurus semitaeniatus
Thecadactylus rapicauda
Borikenophis portoricensis anegadae

Necker Island (18.55° /- 64.35°) (0.30 km^2)
Eleutherodactylus antillensis (I)
Osteopilus septentrionalis (W)
Centrochelys sulcata (I)
Chelonoidis carbonaria (I)
Ameiva exsul
Anolis cristatellus wileyae
Anolis pulchellus
Anolis stratulus
Cyclura pinguis (I)
Sphaerodactylus macrolepis macrolepis
Spondylurus semitaeniatus
Thecadactylus rapicauda
Borikenophis portoricensis anegadae
Typhlops naugus

Prickly Pear Island (18.50° /- 64.37°) (0.70 km^2)
Ameiva exsul
Anolis cristatellus wileyae

| | |
|--|--|
| <i>Anolis stratulus</i> | <i>Osteopilus septentrionalis</i> (I) |
| <i>Sphaerodactylus macrolepis macrolepis</i> | <i>Lithobates catesbeianus</i> (W) |
| <i>Spondylurus</i> sp. | <i>Lithobates pipiens</i> (W) |
| <i>Typhlops naugus</i> | <i>Rhinella marina</i> (I) |
| Round Rock (18.40°/-64.45°) (0.08 km ²) | <i>Chelonoidis carbonaria</i> (I?) |
| <i>Ameiva exsul</i> | <i>Trachemys scripta elegans</i> (I) |
| <i>Anolis cristatellus wileyae</i> | <i>Ameiva exsul</i> (I) |
| <i>Sphaerodactylus macrolepis macrolepis</i> | <i>Ameiva polops</i> (PX) |
| <i>Spondylurus semitaeniatus</i> | <i>Anolis acutus</i> |
| Saba Rock (18.50°/-64.45°) (0.002 km ²) | <i>Capitellum parvicruzae</i> (PE) |
| <i>Anolis cristatellus wileyae</i> | <i>Hemidactylus mabouia</i> (I?) |
| <i>Sphaerodactylus macrolepis macrolepis</i> | <i>Iguana iguana</i> (I) |
| Salt Island (18.38°/-64.52°) (0.78 km ²) | <i>Sphaerodactylus beattyi beattyi</i> |
| <i>Ameiva exsul</i> | <i>Sphaerodactylus beattyi seamanii</i> |
| <i>Anolis cristatellus wileyae</i> | <i>Sphaerodactylus macrolepis macrolepis</i> |
| <i>Anolis stratulus</i> | <i>Spondylurus magnacruzae</i> (PE) |
| <i>Hemidactylus mabouia</i> (I?) | <i>Thecadactylus rapicauda</i> |
| <i>Sphaerodactylus macrolepis macrolepis</i> | <i>Borikenophis portoricensis</i> (W) |
| <i>Spondylurus semitaeniatus</i> | <i>Borikenophis sancticrucis</i> (PE) |
| <i>Spondylurus sloanii</i> | <i>Pantherophis guttatus</i> (W) |
| <i>Borikenophis portoricensis richardi</i> | <i>Ramphotyphlops braminus</i> (I) |
| South Cockroach (= Big Cockroach = Cockroach Island) (18.48°/-64.45°) (0.004 km ²) | <i>Thamnophis</i> sp. (W) |
| <i>Anolis cristatellus wileyae</i> | <i>Typhlops richardii</i> |
| <i>Sphaerodactylus macrolepis macrolepis</i> | Buck Island (17.79°/-64.62°) (0.73 km ²) |
| West Dog (18.48°/-64.47°) (0.13 km ²) | <i>Ameiva polops</i> (I) |
| <i>Anolis cristatellus wileyae</i> | <i>Anolis acutus</i> |
| <i>Sphaerodactylus macrolepis macrolepis</i> | <i>Sphaerodactylus beattyi beattyi</i> |
| West Seal Dog (18.50°/-64.43°) (0.02 km ²) | <i>Sphaerodactylus macrolepis macrolepis</i> |
| <i>Anolis cristatellus wileyae</i> | Green Cay (17.77°/-64.67°) (0.05 km ²) |
| <i>Sphaerodactylus macrolepis macrolepis</i> | <i>Ameiva polops</i> |
| ANEGADA (18.75°/-64.33°) (38.72 km ²) | <i>Anolis acutus</i> |
| <i>Leptodactylus albobilabris</i> | <i>Sphaerodactylus beattyi beattyi</i> |
| <i>Ameiva exsul</i> | <i>Sphaerodactylus macrolepis macrolepis</i> |
| <i>Anolis cristatellus wileyae</i> | <i>Spondylurus magnacruzae</i> (PE) |
| <i>Anolis pulchellus</i> | <i>Borikenophis sancticrucis</i> (PE) |
| <i>Anolis stratulus</i> | Protestant Cay (17.75°/-64.70°) (0.02 km ²) |
| <i>Cyclura pinguis</i> | <i>Ameiva polops</i> |
| <i>Hemidactylus mabouia</i> (I?) | <i>Anolis acutus</i> |
| <i>Sphaerodactylus macrolepis macrolepis</i> | <i>Sphaerodactylus macrolepis macrolepis</i> |
| <i>Spondylurus anegadae</i> | Ruth Island (17.68°/-64.76°) (0.08 km ²) |
| <i>Thecadactylus rapicauda</i> | <i>Ameiva polops</i> (I) |
| <i>Borikenophis portoricensis anegadae</i> | |
| <i>Magliophis exiguum exiguum</i> | |
| <i>Typhlops catapontus</i> | |

ST. CROIX BANK

| |
|---|
| ST. CROIX (17.74°/-64.74°) (211.78 km ²) |
| <i>Eleutherodactylus antillensis</i> (I?) |
| <i>Eleutherodactylus coqui</i> (I) |
| <i>Eleutherodactylus latus</i> |
| <i>Eleutherodactylus schwartzi</i> (I?) |
| <i>Leptodactylus albobilabris</i> |

LITERATURE CITED

- Alayo, P. 1955. Lista de los Anfibios de Cuba. Museo Charles Ramsden, Universidad de Oriente, Santiago de Cuba, 11 p.
- Barbour, T. 1914. A contribution to the zoogeography of the West Indies, with especial reference to amphibians and reptiles. Memoirs of the Museum of Comparative Zoölogy 44:205–359.
- Barbour, T. 1930. A list of Antillean reptiles and amphibians. *Zoologica* (N.Y.) 11:61–116.
- Barbour, T. 1935. A second list of Antillean reptiles and amphibians. *Zoologica* (N.Y.) 19:77–141.
- Barbour, T. 1937. Third list of Antillean reptiles and amphibians. *Bulletin of the Museum of Comparative Zoölogy* 82:77–166.
- Barbour T., & C. T. Ramsden. 1919. The herpetology of Cuba. *Memoirs of the Museum of Comparative Zoölogy* 47:71–213.
- Barbour, T., & B. Shreve. 1935. Concerning some Bahamian reptiles, with notes on the fauna. *Proceedings of the Boston Society of Natural History* 40:347–366.
- Breuil, M. 2002. Histoire naturelle des amphibiens et reptiles terrestres de l'archipel Guadeloupéen. Guadeloupe, Saint-Martin, Saint Barthélémy. Patrimoines Naturels, Paris 54:339 pp.
- Breuil, M. 2011. The terrestrial herpetofauna of Martinique: Past, present, future. Pp. 311–338 in A. Hailey, B. S. Wilson, & J. A. Horrocks, eds. *Conservation of Caribbean Island Herpetofaunas. Volume 2: Regional Accounts of the West Indies*. Brill, Leiden, The Netherlands.
- Bryan, J. J., G. P. Gerber, M. E. Welch, & C. L. Stephen. 2007. Re-evaluating the taxonomic status of the Booby Cay Iguana, *Cyclura carinata bartschi*. *Copeia* 2007:734–739.
- Buide, M. S. 1967. Lista de los anfibios y reptiles de Cuba. *Torreia* 1:1–60.
- Burns, J. K., C. A. Cunningham, R. A. Dupuis, M. N. Trask, J. S. Tulloch, R. Powell, J. S. Parmerlee, Jr., K. L. Kopecky, & M. L. Jolley. 1992. Lizards of the Cayos Siete Hermanos, Dominican Republic, Hispaniola. *Bulletin Chicago Herpetological Society* 27:225–232.
- Carew, J. L., & J. E. Mylroie. 1997. Geology of the Bahamas. Pp. 91–139 in H. L. Vacher & T. M. Quinn, eds. *Geology and Hydrogeology of Carbonate Islands. Developments in Sedimentology Series 54*. Elsevier Science, Amsterdam, 948 p.
- Carlson, L. A. 1999. Aftermath of a feast: Human colonization of the southern Bahama Archipelago and its effects on the indigenous fauna. Ph.D. dissertation, Department of Anthropology, University of Florida, Gainesville, 558 p.
- Censky, E. J. 1988. *Geochelone carbonaria* (Reptilia: Testudines) in the West Indies. *Florida Scientist* 50:108–114.
- Corke, D. 1992. The status and conservation needs of the terrestrial herpetofauna of the Windward Islands (West Indies). *Biological Conservation* 62:47–58.
- Crombie, R. I. 1999. Jamaica. Pp. 63–92 in B.I. Crother, ed. *Caribbean Amphibians and Reptiles*. Academic Press, San Diego, California.
- Daltry, J. C. 2011. An introduction to the herpetofauna of Antigua, Barbuda and Redonda, with some conservation recommendations. Pp. 17–51 in A. Hailey, B. S. Wilson, & J. A. Horrocks, eds. *Conservation of Caribbean Island Herpetofaunas. Volume 2: Regional Accounts of the West Indies*. Brill, Leiden, The Netherlands.
- Daudin, J., & M. de Silva. 2011. The herpetofauna of Grenada and the Grenada Grenadines: Conservation concerns. Pp. 259–271 in A. Hailey, B. S. Wilson, & J. A. Horrocks, eds. *Conservation of Caribbean Island Herpetofaunas. Volume 2: Regional Accounts of the West Indies*. Brill, Leiden, The Netherlands.
- Defense Mapping Agency Topographic Center. 1973. *Haiti. Official Standard Names Gazetteer*. 2nd ed. U.S. Department of the Interior, Washington, DC.
- de la Sagra, R. 1838–1861. *Historia física, política y natural de la Isla de Cuba*. Librería de Arthus Bertrand, Librero de la Sociedad de Geografía, Paris, 242 p.

- Díaz, L. M., & A. Cádiz. 2008. Guía taxonómica de los anfibios de Cuba. *Abc Taxa* 4:294 pp.
- Dunn, E. R. 1936. Notes on American *mabuyas*. *Proceedings of the Academy of Natural Sciences of Philadelphia* 87:533–557.
- Echternacht, A. C., F. J. Burton, & J. M. Blumenthal. 2011. The amphibians and reptiles of the Cayman Islands: Conservation issues in the face of invasions. Pp. 129–147 in A. Hailey, B. S. Wilson, & J. A. Horrocks, eds. *Conservation of Caribbean Island Herpetofaunas. Volume 2: Regional Accounts of the West Indies*. Brill, Leiden, The Netherlands.
- Estrada, A. R. 1992. Lista de anfibios y reptiles de Cayo Saetía, costa N de Mayarí, Holguín, Cuba. *Comunicaciones Cortas de Zoológica*:14–15.
- Estrada, A. R. 1993a. Anfibios y reptiles de Cayo Coco, Archipiélago de Sabana Camagüey, Cuba. *Poeyana* 432:1–21.
- Estrada, A. R. 1993b. Herpetofauna del Archipiélago de los Canarreos, Cuba. *Poeyana* 431:1–19.
- Estrada, A. R. 1994. Herpetofauna de la Cuenca Banao-Higuanojo, Sancti Spíritus, Cuba. *Revista de la Academia Colombiana de Ciencias* 19:353–360.
- Estrada, A. R., & J. Novo. 1984. Reptiles y aves de Cayo Inés de Soto, Archipiélago de los Colorados, Pinar del Río, Cuba. *Miscelánea Zoológica* 23:1.
- Estrada, A. R., & R. Rodríguez. 1985. Lista de vertebrados terrestres de Cayo Campos, Archipiélago de los Canarreos, Cuba. *Miscelánea Zoológica* 28:1–2.
- Estrada, A. R., J. Novo, & L. V. Moreno. 1987. Datos sobre una puesta de *Ameiva auberti* Cocteau (Sauria: Teiidae). *Miscelánea Zoológica* 30:2–3.
- Fairbanks, R. G. 1989. A 17,000-year glacio-eustatic sea level record: Influence of glacial melting rates on the Younger Dryas event and deep-ocean circulation. *Nature* 342:637–642.
- Fields, A., & J. A. Horrocks. 2011. The herpetofauna of Barbados: Anthropogenic impacts and conservation status. Pp. 89–104 in A. Hailey, B. S. Wilson, & J. A. Horrocks, eds. *Conservation of Caribbean Island Herpetofaunas. Volume 2: Regional Accounts of the West Indies*. Brill, Leiden, The Netherlands.
- Franz, R., C. K. Dodd, & S. D. Buckner. 1996. A review of herpetology of the Bahamian Archipelago. *Bahamas Journal of Science* 3(3):22–30.
- Franz, R., C. K. Dodd, & D. W. Buden. 1993. Distributional records of amphibians and reptiles from the Exuma Islands, Bahamas, including the first reports of a freshwater turtle and an introduced gecko. *Caribbean Journal of Science* 29:165–173.
- Franz, R., & S. Franz. 2009. A new fossil tortoise in the genus *Chelonoidis* (Testudinidae) from the northern Bahamas, with an osteological assessment of other Neotropical tortoises. *Bulletin of the Florida Museum of Natural History* 49:1–44.
- Garrido, O. H. 1973a. Anfibios, reptiles y aves de Cayo Real (Cayos de San Felipe), Cuba. *Poeyana* 119:1–49.
- Garrido, O. H. 1973b. Distribución y variación de *Anolis homolechis* Cope (Lacertilia: Iguanidae) en Cuba. *Poeyana* 120:1–64.
- Garrido, O. H. 1976. Aves y reptiles de Cayo Coco, Cuba. *Miscelánea Zoológica* 3:1–4.
- Garrido, O. H. 1980. Los vertebrados terrestres de la Península de Zapata. *Poeyana* 203:1–49.
- Garrido, O. H., & M. L. Jaume. 1984. Catálogo descriptivo de los anfibios y reptiles de Cuba. Doñana, *Acta Vertebrata* 11:5–128.
- Garrido, O. H., A. R. Estrada, & A. Llanes. 1986. Anfibios, reptiles y aves de Cayo Guajaba, Archipiélago de Sabana-Camagüey, Cuba. *Poeyana* 328:1–34.
- Gerber, G. P. 1998. Management plan for the protection of the iguana. Pp. C1–C37 in *Strategic Environmental Assessment for Big Ambergris Cay, Turks and Caicos Islands, British West Indies*. Coastal Systems International, Coral Gables, Florida.
- Glor, R. E., & R. Laport. In press. Are subspecies of *Anolis* lizards (*A. distichus*) that differ in dewlap color and pattern also genetically distinct? A mitochondrial analysis. *Molecular Phylogenetics and Evolution* (doi:10.1016/j.ympev.2010.11.004).

- Grant, C. 1937. Herpetological notes with new species from the American and British Virgin Islands, 1936. *Journal of Agriculture, University of Puerto Rico* 21:503–522.
- Grant, C. 1941 [“1940”]. The herpetology of the Cayman Islands. *Bulletin of the Institute of Jamaica, Science Series* 2:65 pp.
- Greene, B. T., D. T. Yorks, J. S. Parmerlee, Jr., R. Powell, & R. W. Henderson. 2002. Discovery of *Anolis sagrei* in Grenada with comments on its potential impact on native anoles. *Caribbean Journal of Science* 38:270–272.
- Gundlach, J. 1867. Revista y catálogo de los reptiles cubanos. *Repertorio Físico Natural de la Isla de Cuba* 2:102–119.
- Gundlach, J. 1880. Contribución a la Herpetología Cubana. G. Montiel, La Habana, 99 p.
- Heatwole, H., & F. MacKenzie. 1967. Herpetogeography of Puerto Rico. IV. Paleogeography, faunal similarity and endemism. *Evolution* 21:429–439.
- Heatwole, H., R. Levins, & M. D. Byer. 1981. Biogeography of the Puerto Rican Bank. *Atoll Research Bulletin* 251:1–55.
- Hedges, S. B. 1999. Distribution patterns of amphibians in the West Indies. Pp. 211–254 in W. E. Duellman, ed. *Patterns of Distribution of Amphibians: A Global Perspective*. Johns Hopkins University Press. Baltimore, Maryland.
- Hedges, S. B. 2012. Caribherp: West Indian Amphibians and Reptiles (<http://caribherp.org>). Pennsylvania State University, University Park.
- Hedges, S. B., & C. E. Conn. 2012. A new skink fauna from Caribbean islands (Squamata, Scincidae, Mabuini). *Zootaxa* 3228:1–244.
- Henderson, R. W., & C. S. Berg. 2011. The herpetofauna of Grenada and the Grenada Grenadines: Conservation concerns. Pp. 239–258 in A. Hailey, B. S. Wilson, & J. A. Horrocks, eds. *Conservation of Caribbean Island Herpetofaunas. Volume 2: Regional Accounts of the West Indies*. Brill, Leiden, The Netherlands.
- Henderson, R. W., & R. Powell. 2001. Responses by the West Indian herpetofauna to human-influenced resources. *Caribbean Journal of Science* 37:41–54.
- Henderson, R. W., & R. Powell. 2005. Thomas Barbour and the *Utarwana* voyages (1929–1934) in the West Indies. *Bonner Zoologische Beiträge* 52:297–309.
- Henderson, R. W., & R. Powell. 2009. *Natural History of West Indian Reptiles and Amphibians*. University Press of Florida, Gainesville, 496 p.
- Hodge, K. V. D., E. J. Censky, & R. Powell. 2003. *The Reptiles and Amphibians of Anguilla, British West Indies*. Anguilla National Trust, The Valley, Anguilla, British West Indies, 72 p.
- Hodge, K. V. D., R. Powell, & E. J. Censky. 2011. Conserving the herpetofauna of Anguilla. Pp. 3–15 in A. Hailey, B. S. Wilson, & J. A. Horrocks, eds. *Conservation of Caribbean Island Herpetofaunas. Volume 2: Regional Accounts of the West Indies*. Brill, Leiden, The Netherlands.
- IUCN (International Union for the Conservation of Nature and Natural Resources). 2011. 2011 IUCN Red List of Threatened Species. <www.iucnredlist.org>
- Iverson, J. B. 1978. The impact of feral cats and dogs on populations of the West Indian rock iguana, *Cyclura carinata*. *Biological Conservation* 14:63–73.
- Jaume, M. L. 1966. Catálogo de la Fauna de Cuba. XV. Catálogo de los anfibios de Cuba. Museo “Felipe Poey” de la Academia de Ciencias de la República de Cuba, Trabajos de Divulgación 35:1–20.
- Johnson, S. 2011. A non-native skink on Grand Bahama. *Reptiles & Amphibians* 18:247–248.
- Kardong, K. V. 1980. Jaw musculature of the West Indian snake *Alsophis cantherigerus brooksi* (Colubridae, Reptilia). *Breviora* 463:1–26.
- Knapp, C. R., J. B. Iverson, S. D. Buckner, & S. V. Cant. 2011. Conservation of amphibians and reptiles in the Bahamas. Pp. 53–87 in A. Hailey, B. S. Wilson, & J. A. Horrocks, eds. *Conservation of Caribbean Island Herpetofaunas. Volume 2: Regional Accounts of the West Indies*. Brill, Leiden, The Netherlands.
- Köhler, G., & M. Vesely. 2011. A new species of *Thecadactylus* from Sint Maarten, Lesser Antilles (Reptilia, Squamata, Gekkonidae).

- ZooKeys 118:97–107.
- Kraus, F. 2009. Alien Reptiles and Amphibians: A Scientific Compendium and Analysis. Invading Nature: Springer Series in Invasion Biology 4. Springer, New York, 563 p.
- Lazell, J. 1983. Biogeography of the herpetofauna of the British Virgin Islands, with description of a new anole (Sauria: Iguanidae). Pp. 99–117 in A. G. J. Rhodin & K. Miyata, eds. Advances in Herpetology and Evolutionary Biology. Museum of Comparative Zoology, Cambridge, Massachusetts.
- Lazell, J. 1996. Careening Island and the Goat Islands: Evidence for the arid-insular invasion wave theory of dichopatric speciation in Jamaica. Pp. 195–205 in R. Powell & R. W. Henderson, eds. Contributions to West Indian Herpetology: A Tribute to Albert Schwartz. Society for the Study of Amphibians and Reptiles Contributions to Herpetology, vol. 12. Ithaca, New York.
- Lazell, J. 2005. Island: Fact and Theory in Nature. University of California Press, Berkeley, 382 p.
- Lee, D. S. 2004. Additional reptiles and amphibians introduced to the Bahamas: A growing conservation concern. Bulletin of the Chicago Herpetological Society 39:161–164 (same text appears in the Bahamas Journal of Science 12(2):2–6).
- Lever, C. 2003. Naturalized Reptiles and Amphibians of the World. Oxford University Press, New York, 318 p.
- Lorvelec, O., M. Pascal, C. Pavis, & P. Feldmann. 2011. Amphibians and reptiles of the French West Indies: Inventory, threats and conservation. Pp. 205–237 in A. Hailey, B. S. Wilson, & J. A. Horrocks, eds. Conservation of Caribbean Island Herpetofaunas. Volume 2: Regional Accounts of the West Indies. Brill, Leiden, The Netherlands.
- MacLean, W. P. 1982. Reptiles and Amphibians of the Virgin Islands. Macmillan, London, 54 p.
- MacLean, W. P., R. Kellner, & H. Dennis. 1977. Island lists of West Indian amphibians and reptiles. Smithsonian Herpetological Information Service(40:1–47).
- Malhotra, A., & R.S. Thorpe. 1999. Reptiles and Amphibians of the Eastern Caribbean. MacMillan Education Ltd., London, 134 p.
- Malhotra, A., R. S. Thorpe, E. Hypolite, & A. James. 2011. A report on the status of the herpetofauna of the Commonwealth of Dominica, West Indies. Pp. 149–166 in A. Hailey, B. S. Wilson, & J. A. Horrocks, eds. Conservation of Caribbean Island Herpetofaunas. Volume 2: Regional Accounts of the West Indies. Brill, Leiden, The Netherlands.
- Mayer, G. C. 1989. Deterministic aspects of community structure in West Indian amphibians and reptiles. Ph.D. thesis, Harvard University, Cambridge, Massachusetts, 294 p.
- Mayer, G. C., & J. Lazell. 2000. A new species of *Mabuya* (Sauria: Scincidae) from the British Virgin Islands. Proceedings of the Biological Society of Washington 113:871–886.
- McGuire, J. W. 1925. Geographic Dictionary of the Virgin Islands of the United States. U.S. Coast and Geodetic Survey Special Publication 103:1–211.
- Myers, N., R. A. Mittermeier, C. G. Mittermeier, G. A. B. da Fonseca, & J. Kent. 2000. Biodiversity hotspots for conservation priorities. Nature 403:853–858.
- Newsom, L. A., & E. S. Wing 2004. On Land and Sea: Native American Uses of Biological Resources in the West Indies. University of Alabama Press, Tuscaloosa, 323 p.
- Ng, J., & R. E. Glor. 2011. Genetic differentiation among populations of a Hispaniolan trunk anole that exhibit geographical variation in dewlap colour. Molecular Ecology 20:4302–4317.
- Novo, J., A. R. Estrada, & L. V. Moreno 1987. Adiciones a la fauna de anfibios de la Península de Guanahacabibes, Cuba. Miscelánea Zoológica 36:3–4.
- Ovaska, K. E., J. Caldbeck, & J. Lazell. 2000. New records and distributional and ecological notes of leptodactylid frogs, *Leptodactylus* and *Eleutherodactylus*, from the British Virgin Islands. Breviora 508:1–25.
- Perry, G., & G. P. Gerber. 2011. Conservation of amphibians and reptiles in the British Virgin Islands. Pp. 105–127 in A. Hailey, B. S. Wilson, & J. A. Horrocks, eds. Conservation of Caribbean Island Herpetofaunas. Volume 2:

- Regional Accounts of the West Indies. Brill, Leiden, The Netherlands.
- Philibosian, R., & J. A. Yntema. 1976. Records and status of some reptiles and amphibians in the Virgin Islands. I. 1968–1975. *Herpetologica* 32:81–85.
- Philibosian, R., & J. A. Yntema. 1978. Records and status of some reptiles and amphibians in the Virgin Islands. II. 1975–1976. *Herpetologica* 34:47–51.
- Platenberg, R. J. 2007. Impacts of introduced species on an island ecosystem: Non-native reptiles and amphibians in the U.S. Virgin Islands. Pp. 168–174 in G. W. Witmer, W. C. Pitt, & K. A. Fagerstone, eds. *Managing Vertebrate Invasive Species: Proceedings of an International Symposium*. National Wildlife Research Center, Fort Collins, Colorado.
- Platenberg, R. J., & R. H. Boulon, Jr. 2011. Conservation status of reptiles and amphibians in the U.S. Virgin Islands. Pp. 407–428 in A. Hailey, B. S. Wilson, & J. A. Horrocks, eds. *Conservation of Caribbean Island Herpetofaunas. Volume 2: Regional Accounts of the West Indies*. Brill, Leiden, The Netherlands.
- Powell, R. 1999. Herpetology of Navassa Island, West Indies. *Caribbean Journal of Science* 35:1–13.
- Powell, R. 2011. Conservation of the herpetofauna on the Dutch Windward Islands: St. Eustatius, Saba, and St. Maarten. Pp. 189–204 in A. Hailey, B. S. Wilson, & J. A. Horrocks, eds. *Conservation of Caribbean Island Herpetofaunas. Volume 2: Regional Accounts of the West Indies*. Brill, Leiden, The Netherlands.
- Powell, R., & R. W. Henderson. 1999. Addenda to the checklist of West Indian amphibians and reptiles. *Herpetological Review* 30:137–139.
- Powell, R., & R. W. Henderson. 2003. A second set of addenda to the checklist of West Indian amphibians and reptiles. *Herpetological Review* 34:341–345.
- Powell, R., & R. W. Henderson. 2011. The St. Vincent (Lesser Antilles) herpetofauna: Conservation concerns. Pp. 359–376 in A. Hailey, B. S. Wilson, & J. A. Horrocks, eds. *Conservation of Caribbean Island Herpetofaunas. Volume 2: Regional Accounts of the West Indies*. Brill, Leiden, The Netherlands.
- Powell, R., & S. J. Incháustegui. 2011. Conservation of the herpetofauna of the Dominican Republic. Pp. 167–188 in A. Hailey, B. S. Wilson, & J. A. Horrocks, eds. *Conservation of Caribbean Island Herpetofaunas. Volume 2: Regional Accounts of the West Indies*. Brill, Leiden, The Netherlands.
- Powell, R., R. W. Henderson, K. Adler, & H. A. Dundee. 1996. An annotated checklist of West Indian amphibians and reptiles. Pp. 51–93 in R. Powell & R. W. Henderson, eds. *Contributions to West Indian Herpetology: A Tribute to Albert Schwartz*. Society for the Study of Amphibians and Reptiles Contributions to Herpetology, vol. 12. Ithaca, New York.
- Powell, R., J. A. Ottenwalder, & S. J. Incháustegui. 1999. The Hispaniolan herpetofauna: diversity, endemism, and historical perspectives, with comments on Navassa Island. Pp. 93–168 in B.I. Crother, ed. *Caribbean Amphibians and Reptiles*. Academic Press, San Diego, California.
- Powell, R., R. W. Henderson, & J. S. Parmerlee Jr. 2005. *Reptiles and Amphibians of the Dutch Caribbean: St. Eustatius, Saba, and St. Maarten*. St. Eustatius National Parks Foundation, Gallows Bay, St. Eustatius, Netherlands Antilles, 192 p.
- Powell, R., R. W. Henderson, M. C. Farmer, M. Breuil, A. C. Echternacht, G. van Buurt, C. M. Romagosa, & G. Perry. 2011. Introduced amphibians and reptiles in the greater Caribbean: Patterns and conservation implications. Pp. 63–143 in A. Hailey, B. S. Wilson, & J. A. Horrocks, eds. *Conservation of Caribbean Island Herpetofaunas. Volume 1: Conservation Biology and the Wider Caribbean*. Brill, Leiden, The Netherlands.
- Pregill, G. K., & B. I. Crother. 1999. Ecological and historical biogeography. Pp. 335–356 in B. I. Crother, ed. *Caribbean Amphibians and Reptiles*. Academic Press, San Diego.
- Pregill, G. K., D. W. Steadman, & D. R. Watters. 1994. Late Quaternary vertebrate faunas of the Lesser Antilles: historical components

- of Caribbean biogeography. *Bulletin of the Carnegie Museum of Natural History* 30:1–51.
- Reynolds, R. G. 2010. *Typhlops platycephalus* (Puerto Rican White-tailed Blindsnake), distribution. *Caribbean Herpetology* 1:15.
- Reynolds, R. G. 2011a. Status, conservation, and introduction of amphibians and reptiles in the Turks and Caicos Islands, British West Indies. Pp. 377–406 in A. Hailey, B. S. Wilson, & J. A. Horrocks, eds. *Conservation of Caribbean Island Herpetofaunas. Volume 2: Regional Accounts of the West Indies*. Brill, Leiden, The Netherlands.
- Reynolds, R. G. 2011b. Islands, metapopulations, and archipelagos: Genetic equilibrium and non-equilibrium dynamics of structured populations in the context of conservation. Ph.D. dissertation, Department of Ecology and Evolutionary Biology, University of Tennessee, Knoxville, 185 p.
- Reynolds, R. G. 2012. *Epicrates chrysogaster*. Catalogue of American Amphibians and Reptiles 898:1–5.
- Reynolds, R. G., & M. L. Niemiller. 2009. *Hemidactylus mabouia* (Wood Slave). Geographic distribution. *Herpetological Review* 40:452.
- Reynolds, R. G., & M. L. Niemiller. 2010a. Island invaders: Introduced reptiles and amphibians of the Turks and Caicos Islands. *Reptiles & Amphibians* 17:117–121.
- Reynolds, R. G., & M. L. Niemiller. 2010b. *Epicrates chrysogaster* (Southern Bahamas Boa), distribution. *Caribbean Herpetology* 1:14.
- Reynolds, R. G., G. P. Gerber, & J. Burgess. 2010. *Tropidophis greenwayi greenwayi* (Big Ambergris Dwarf Boa). Geographic distribution. *Herpetological Review* 41:520.
- Reynolds, R. G., G. P. Gerber, & B. M. Fitzpatrick. 2011. Unexpected shallow genetic divergence in Turks Island Boas (*Epicrates c. chrysogaster*) reveals single evolutionary significant unit for conservation. *Herpetologica* 67:477–486.
- Reynolds, R. G., M. L. Niemiller, & B. M. Fitzpatrick. 2012. Genetic analysis of an endemic archipelagic lizard reveals sympatric cryptic lineages and taxonomic discordance. *Conservation Genetics (online first, doi: 10.1007/s10592-012-0344-z)*.
- Rivero, J. A. 1978. Los Anfibios y Reptiles de Puerto Rico. *The Amphibians and Reptiles of Puerto Rico*. Editorial de la Universidad de Puerto Rico, Río Piedras, 152 p. (vol. 1), 148 p. (vol. 2), 63 p. (plates).
- Rivero, J. A. 1998. Los Anfibios y Reptiles de Puerto Rico, segunda edición revisada. *The Amphibians and Reptiles of Puerto Rico*, second edition revised. Editorial de la Universidad de Puerto Rico, Río Piedras, 510 p.
- Rodríguez Schettino, L. (ed.). 1999. *The Iguanid Lizards of Cuba*. University Press of Florida, Gainesville, 428 p.
- Rodríguez Schettino, L. (ed.). 2003. *Anfibios y Reptiles de Cuba*. UPC Print, Vaasa, Finland, 169 p.
- Schmidt, K. P. 1928. Amphibians and land reptiles of Porto Rico, with a list of those reported from the Virgin Islands. *Scientific Survey of Porto Rico and the Virgin Islands* 10:1–160.
- Schwartz, A., & R. W. Henderson. 1988. West Indian amphibians and reptiles: A check-list. *Milwaukee Public Museum Contributions in Biology and Geology* (74):1–264.
- Schwartz, A., & R. W. Henderson. 1991. *Amphibians and Reptiles of the West Indies: Descriptions, Distributions, and Natural History*. University of Florida Press, Gainesville, 720 p.
- Schwartz, A., & R. Thomas. 1975. A check-list of West Indian amphibians and reptiles. *Carnegie Museum of Natural History Special Publication* 1:1–216.
- Seidel, M. E., & R. Franz. 1994. Amphibians and reptiles (exclusive of marine turtles) of the Cayman Islands. Pp. 407–433 in M. A. Brunt & J. E. Davies, eds. *The Cayman Islands: Natural History and Biogeography*. Kluwer Academic Publishers, Dordrecht, The Netherlands.
- Steadman, D. W., R. Franz, G. S. Morgan, N. A. Albury, B. Kakuk, K. Broad, S. E. Franz, K. Tinker, M. P. Pateman, T. A. Lott, D. M. Jarzen, & D. L. Dilcher. 2007. Exceptionally well preserved late Quaternary plant and vertebrate fossils from a blue hole on Abaco, The Bahamas. *Proceedings of the National Academy of Sciences* 104:19897–19902.
- Stejneger, L. 1904. The herpetology of Porto Rico. *Report of the United States National Museum* 1902:549–724.

- Stuart, S. N., J. S. Chanson, N. A. Cox, B. E. Young, A. S. L. Rodrigues, D. L. Fischman, & R. W. Waller. 2004. Status and trends of amphibian declines and extinctions worldwide. *Science* 306:1783–1786.
- Wetmore, A. 1927. The birds of Porto Rico and the Virgin Islands. *Scientific Survey of Porto Rico and the Virgin Islands* 9:243–571.
- Wilson, B. S. 2011. Conservation of Jamaican amphibians and reptiles. Pp. 273–310 *in* A. Hailey, B. S. Wilson, & J. A. Horrocks, eds. *Conservation of Caribbean Island Herpetofaunas. Volume 2: Regional Accounts of the West Indies*. Brill, Leiden, The Netherlands.
- Wing, E. S., & S. J. Scudder 1983. Animal exploitation by people living on a tropical marine edge. Pp. 197–210 *in* C. Grigson & J. Clutton-Brock, eds. *Animals and Archeology: 2. Shell Middens, Fishes and Birds*. British Archaeological Reports International Series 183. Archaeopress, Oxford, UK.
- Yeska, M. L., R. Powell, & J. S. Parmerlee Jr. 2000. Lizards of Cayo Pisaje, Dominican Republic, Hispaniola. *Herpetological Review* 31:18–20.